AUTHOR INDEX

Alder, F. E., *see* Hughes, G. P.

Badreldin, A. L., *see* Hafez, E. S. E.
Blackett, R. D., dry-matter in fodder-beet, 202–5.
Bolton, A., tobacco-seed production, 241–3.
Bunting, A. H., groundnut varieties for Tanganyika, 39–44; classification of groundnuts, 158–69.

Carlisle, Elizabeth, influence of fertilizers on secretion of nectar by red clover, 126–30.
Castle, M. E., fodder-beet for fattening ducks, 251–6.
Child, R., fertilizers for Napier grass, 220–5.
Christidis, B. G., rotation experiments with cotton, 49–54.
Clapperton, J. L., *see* Thomas, Brynmor.
Cooke, F. W., performance of fertilizer distributors, 244–50.

Davies, R. O., *see* Henderson, J. L.

Foot, A. S., marrowstem kale silage, 109–12.

Gokhale, N. G., decrease in yield of tea on ceasing to manure, 96–100.
Goodchild, N. A., *see* Child, R.

Hafez, E. S. E., hair coat in Bovinae, 34–38.
Hancock, J., effect of tropical climate on European-type cattle. I. Growth, 55–74.
Hassan, H. A., *see* Sirry, I.
Hauser, G., comparison of Afghan ploughing with modern implements and methods, 75–80.
Henderson, J. L., yield and composition of cereal-legume mixtures at different stages of growth, 131–44.
Hewitt, C. W., leaf analysis as guide to banana nutrition, 11–16.
Holmes, W. B., *see* Thomas, Brynmor.
Hughes, G. P., overwintering of beef stores, 145–57.

Jones, K. Wilson, witchweed control. II. Physiological strains of *Striga hermontheca*, 206–13.
Josephson, L. M., cytoplasmic male sterility in production of hybrid maize, 1–10.

King, J. O. L., *see* Castle, M. E.
Knight, R., introductions to West Africa of *Theobroma cacao* and related species. I. Review of the first ten years, 113–25.

Lea, J. D., *see* Bunting, A. H.

McLaughlin, E. A., cattle of the Fung, a variant of the Shorthorn Zebu of N. Sudan, 188–201.
Magee, A. I., tuber bruising and potato-digger operation, 45–48.
Masefield, G. B., nodulation of legumes, 17–24.
Milne, A. H., the humps of East African cattle, 234–9.
Murdoch, J. C., *see* Foot, A. S.

Nutman, F. J., 'lethal yellowing', the 'unknown disease' of coconut palms in Jamaica, 258–67; 'frond drop', an abnormal condition of coconut palms in Jamaica, 268–70.

Oyenuga, V. A., composition and nutritive value of feedingstuffs in Nigeria. I. Roots, tubers, and green leaves, 81–95. II. Concentrated feedingstuffs, 171–87.

Payne, W., *see* Hancock, J.

Redford, R. A., *see* Hughes, G. P.
Roberts, F. M., *see* Nutman, F. J.
Rogers, H. H., *see* Knight, R.
Rowland, S. J., *see* Foot, A. S.
Ryle, Margaret, *see* Carlisle, Elizabeth.

Schimmer, F. C., *see* Selman, I. W.
Selman, I. W., disease incidence and yield of potatoes grown with organic and inorganic fertilizers, 226–33.
Shafei, M. M., *see* Hafez, E. S. E.
Sirry, I., effect of potassium-iodide feeding on milk production of Egyptian buffalo, 214–19.

Thomas, Brynmor, meadow hays from Cockle Park. I. Proximate constituents and digestibility, 25–33. II. Ash constituents, 101–8.
Todd, J. R., *see* Child, R.

SUBJECT INDEX

Banana, nutrition, leaf analysis as guide, 11–16; variation in leaf-composition with age and locality, 12–13; effect of fertilizers on leaf-composition and yield, 13–16; critical levels, 16.
Beef stores, overwintering, 145–57; liveweight increases, after summer grazing, Hereford and Hereford-cross bullocks out- or in-wintered, 148–56; carcass gradings and weights, 155; bullocks out-wintered superior to in-wintered, 156–7.
Beet, fodder, *see* Fodder beet, 202 and 251.
Beet, sugar: yields with modern and native tillage methods in Afghanistan, *see* Tillage.
Bovine hump, East African cattle, 234–9; appearance, 234; structure, 234–5; E. African Shorthorn Zebu, 235–6; Ankole, 236–7; comparison, 237–8.

XUM

Bovines, hair coat, Egyptian buffaloes and cattle, 34–38; structure and distribution of hairs, 36–37; variations with species, breed, sex, body region, age, 36–38.
Buffalo, Egyptian, milk production, effect of potassium-iodide feeding, 214–19; milk yields, lactation period, and composition of milk, 214–17; effect of potassium iodide feeding on milk yield, 216; on lactation period, 216–17; on composition of milk, 217–18; on general condition, 218–19.

Cacao, recent introductions, West Africa. I. Review of first ten years, 113–25; sources of varieties of *Theobroma cacao*, 113–16, and of other Theobroma and Herrania species, 123–4; Upper Amazon population, 114–15; Criollo population, 115–16; Trinitario complex, 116; Lower Amazon, 116; development work: distribution, 116–17; breeding programme, 117–20; yields, 120–1; bean characters, 121–2; future prospects, 125.
Cattle, *see*: beef stores, 145; humps of, 234; hair coat, 34; effect of climate, 55; of Fung (Sudan), 188; *see also* Buffalo, 214.
Cattle, European-type, effect of tropical climate. I. Growth, 55–74; climate, in Fiji and New Zealand, 55–58; identical twins used in experiments, 58–59; growth measurements, 60–69; feed intake and utilization, 69–71; poor nutrition or management probable chief cause of retarded growth in tropics, rather than climate *per se*, 71–73.
Cattle of the Fung, a variant of Shorthorn Zebu of N. Sudan, 188–201; description of region, tribes, and herds, 188–9; Gezira Research Farm: situation, soil, and climate, 189; herd: feeding, 189–90, health, 190, breeding, 190–1, conformation, 191–4, temperament, 194, climatic adaptability, 194–5, breeding performance, 195–6, milk production, 196–8, work performance, 198; possible value for improving milk production in tropics, 198–200.
Cereal-Legume mixtures, yield and composition, at different growth stages, 131–44; oats, peas, and beans grown separately and as a mixture, 131–3; yields and analysis of each crop at different stages of growth, 133–6; comparison of yields of dry matter and crude protein from the four crops, 136–7; curing and harvesting losses, 137–40; yields of animal nutrients in grain and straw less than in immature crops suitable for ensiling, 142–4.
Clover, red, secretion of nectar, influence of fertilizers, 126–30; experimental technique, 126–7; nectar yields with different fertilizers, 127–9; effects of each type of fertilizer, 129–30.
Coconut palms; 'frond drop', an abnormal condition in Jamaica, 268–70; symptoms, 268–9; evidence for pathogenic origin, 269–70.
Coconut palms, 'lethal yellowing', the 'unknown disease' in Jamaica, 257–67; history, 257–8; symptoms, 258; epidemiology, 258–60; pathology, 260–1; observations on nature of disease, 261–2; cytology, 262–3; serological tests, 263; attempts to control, 263–4; field indications, 264–5; evidence supporting virus nature, 265–6.
Cotton, rotation experiments, in Greece, 49–54; rotations compared with continuous cotton, 49–50; yields of seed cotton, 51; earliness, lint percentage and lint length, 52; lack of effect, except on earliness, 53.

Diseases: potatoes, *see* potatoes, disease incidence, &c., 226–33; coconuts, *see* coconut palms, 'lethal yellowing', 257–67, and 'frond drop', 268–70.
Ducks, fattening with fodder-beet, *see* fodder beet, &c., 251–6.

Feedingstuffs in Nigeria, composition and nutritive value. I. Roots, tubers, and green leaves, 81–95; constituents, digestibility, and nutritive value of 38 fodder crops grown at Ibadan, 82–88; results compared, and discussed for each crop, 88–94. II. Concentrated feedingstuffs, 171–87; constituents, digestibility, and nutritive value of 38 concentrates, 171–7; results compared, and discussed for each concentrate, 177–86.
Fertilizer distributors, performance of, 244–50; test technique, 244–7; results with granular and powder fertilizers, 247–8; star-wheel and plate-and-flicker distributors compared, 248–9.
Fodder-beet, dry matter, 202–5; method of sampling, 202–3; variation within varieties and individual roots, 204.
Fodder-beet, for fattening ducks, 251–6; analysis of beet and of meals used in experiments, 252–3; live weight gains and killing-out percentages, 253–4; saving in meal consumption by feeding fodder-beet, 254–5.

Grass, Napier, manuring, at Kericho, Kenya, 220–5; yields with various fertilizers, 221–3; composition of grass and recovery of nutrient elements, 223–4; yield increases caused by nitrogen and phosphate, 225.
Groundnuts, cultivated, classification, 158–69; nature of varieties, 158–9; previous classifications, 159–60; diagnostic characters: branching pattern, 160–2, growth habit, 162, seed dormancy, 162–3, stem, branch, and leaflet characters, 163, flower characters, 164, flowering habit, 163–4, pod characters, 164–5; key to classification of the collection at Tozi, Sudan, 165–9.
Groundnuts: production in Tanganyika, choice of varieties, 39–44; management of variety trials, 39–40; trials, Kongwa area, 40–41, Urambo area, 41–42, Southern Province, 43; disease resistance in relation to varieties, 43–44.

Hay, meadow, Cockle Park plots. I. Proximate constituents and digestibility, 25–33; relation of manuring to composition and digestibility, 27–29; nutritive value inversely related to goodness of manurial treatment, 29–30; relative importance of species and maturity for quality of hay, 30–32. II. Ash constituents, 101–8; differences in content of mineral elements related to fertilizers used, 102–4; effect of manuring on value of hay as source of minerals for livestock, 104–7.

Kale, marrowstem, silage, 109–12; losses: of nutrients in large and small silos, 109–11; not related to seepage, 111–12; not reduced by mixture with hay before ensiling, 111–12.

Leaf analysis, as guide to banana nutrition, *see* banana, nutrition of, 11–16.
Legumes, nodulation, conditions affecting, 17–24; unaffected by proximity of other plants, 18–19; damage to nodules by *Sitonia lineata*, and partial control by insecticides, 19–21; correlation between plant weight and nodule weight, 21–22; life span of nodules, 22.

Maize, hybrid, use of cytoplasmic male sterility in production of, 1–10; cytoplasmic male-sterile inbreds, source of, and methods of use in crossing, 2–3; results of experimental crosses, 3–7; yields of male-sterile double crosses, 7–8; bearing of results on commercial production of hybrid maize seed in S. Africa, 8–10.
Manures and fertilizers, effects of: *see* bananas, 11; clover, red, 126; grass, Napier, 220; hay, 25 and 101; potatoes, 226; tea, 96; tobacco, 240.

Nodulation of legumes, *see* legumes, 17.

Potatoes, bruising, in potato-digger operation, 45–48; factors affecting injury, 45–46; experiments with different speeds and adjustments, 47–48; recommendations, 48.
Potatoes, disease incidence and yield, with organic and inorganic fertilizers, 226–33; meteorological data, 227; crop yields and incidence of disease, 227–31; seasonal yield variations associated with climatic and disease factors, 230–1.

Silage, *see* kale, 109; cereal-legume mixtures, 131.
Sterility, male, cytoplasmic, in production of hybrid maize seed, *see* Maize, 1.

Tea, yield, decrease on ceasing to manure, 96–100; tea-growing in Assam, 96; yields when manured with sulphate of ammonia and after manuring stopped, 97–98; results analysed: equation for calculating fall in yield on ceasing to manure, 98–100.
Tillage, with Afghan plough, compared with modern implements and methods, 75–80; Afghan ploughing and field work, 75–76; wheat, growth and yields, 76–77; sugar beet, growth and yields, 77–79; utilization of irrigation water, 78; modern ploughing significantly better for both wheat and beet, 79–80.
Tobacco-seed production, 240–3; effect of removal of leaves, in S. Rhodesia, on seed-yield and crop value, 241; influence of fertilizers, 241–2; effect of spacing, 242; germination percentage, 242.

Wheat: yields with modern and native tillage methods in Afghanistan, *see* Tillage, 75.
Witchweed control. II. Physiological strains of *Striga hermontheca*, 206–13; factors influencing damage to host plant, 206–7; infestation of millets in Kordofan, Sudan: effect of weed-killers, 207–8; infestation of millets and other cereals and grasses by strains of different origin, on sand, silt, or clay soils, 208–11; evidence for existence of two strains of Striga, 211–13.

XUM